

Direct CP Violation in $B^+ \rightarrow J/\psi K^+$ Decay as Probe for New Physics

Wei-Shu Hou^a, Makiko Nagashima^a, and Andrea Soddu^b

^aDepartment of Physics, National Taiwan University, Taipei, Taiwan 10617, R.O.C.

^bDepartment of Particle Physics, Weizmann Institute of Science, Rehovot 76100, Israel

(Dated: February 2, 2008)

Currently there are New Physics hints in mixing-induced CP violation $\mathcal{S}_{\phi K^0}$, $\mathcal{S}_{\pi^0 K^0} < \mathcal{S}_{J/\psi K^0}$, unequal direct CP violation $\mathcal{A}_{K^+\pi^-} \neq \mathcal{A}_{K^+\pi^0}$, and maybe even in measured $\mathcal{S}_{J/\psi K^0}$ vs prediction from global fit to other data. However, these hints either suffer from experimental uncertainties, or uncertain hadronic corrections, and are not yet unequivocal. Motivated by these hints, however, we point out that a unique probe may be *direct CP* violation in $B \rightarrow J/\psi K$ mode. An asymmetry observed at 1% or higher would indicate New Physics.

PACS numbers: 11.30.Er, 11.30.Hv, 13.25.Hw, 12.60.-i, 12.60.Cn

Unprecedented luminosities have been achieved at the asymmetric energy e^+e^- collider B factories, where altogether close to 10^9 $B\bar{B}$ meson pairs have been collected so far. Besides establishing CP violation (CPV) in the B system, a few hints for possible New Physics (NP) have emerged: the difference between time dependent CPV (TCPV) in charmless $b \rightarrow s\bar{q}q$ modes vs $b \rightarrow c\bar{c}s$ modes; the difference in direct CPV (DCPV) asymmetries between $B^0 \rightarrow K^+\pi^-$ vs $B^+ \rightarrow K^+\pi^0$; and maybe even in the slightly lower value for TCPV in $B^0 \rightarrow J/\psi K^0$ mode compared with the predicted value from fits to other data.

All three hints are not unequivocal, and suffer either in experimental significance, or in theoretical interpretation. In this Letter we point out that DCPV in the $B^+ \rightarrow J/\psi K^+$ mode, $\mathcal{A}_{J/\psi K^+}$, could be at the 1% level or higher, if the above hints are true harbingers of NP. Current measurements give [1] $\mathcal{A}_{J/\psi K^+} = 0.018 \pm 0.043 \pm 0.004$ (CLEO), $-0.026 \pm 0.022 \pm 0.017$ (Belle) and $0.03 \pm 0.015 \pm 0.006$ (BaBar), based on 9.7M, 31.9M and 89M $B\bar{B}$ pairs, respectively. Note that this has not been updated since 2003. The average is [2]

$$\mathcal{A}_{J/\psi K^+} = 0.016 \pm 0.016, \quad (\text{PDG 2005}) \quad (1)$$

where a scale factor has been applied to the error. The nominal error should be of order 0.012 for 131M events, hence a statistical error of 0.003 should be attainable with ~ 2000 M events expected by 2008.

To improve statistics, one could combine with $B^0 \rightarrow J/\psi K^0$ mode. Since Standard Model (SM) expectation is at the sub-percent level, in the next few years $\mathcal{A}_{J/\psi K}$ could be a better probe of NP if seen above 1% level. Such “precision measurement” studies would be good preparation for the future Super B factory, where systematic issues would become a main concern.

Since a few years, a contrast has emerged in mixing-induced TCPV in B decays, i.e. $\mathcal{S}_f$ measurement in a host of CP eigenstates f . The current world average of $\mathcal{S}_f$ in $b \rightarrow c\bar{c}s$ decays is $\sin 2\phi_1/\beta = 0.685 \pm 0.032$ [3], which is dominated by the $B^0 \rightarrow J/\psi K_S$ mode. TCPV measurements in loop dominated $b \rightarrow s\bar{q}q$ processes such as $B^0 \rightarrow \eta' K^0$, ϕK^0 and $\pi^0 K^0$, on the other hand, have persistently given values below $\sin 2\phi_1/\beta$. The current

average in $b \rightarrow s\bar{q}q$ decays is [3] $\mathcal{S}_{s\bar{q}q} = 0.50 \pm 0.06$. The difference $\Delta\mathcal{S} = \mathcal{S}_{s\bar{q}q} - \sin 2\phi_1/\beta$ is only at 2.7σ level. However, $\Delta\mathcal{S}$ has been diminishing experimentally in the past two years, while theoretical interpretation suffers from hadronic uncertainties [4].

DCPV was recently observed [5] in $B^0 \rightarrow K^+\pi^-$ decay, giving $\mathcal{A}_{K^+\pi^-} = -0.108 \pm 0.017$ [3]. However, no indication was seen in charged $B^+ \rightarrow K^+\pi^0$, giving $\mathcal{A}_{K^+\pi^0} = 0.04 \pm 0.04$. The difference with $\mathcal{A}_{K^+\pi^-}$ could be due to an enhancement of color-suppressed amplitude C [6], or due to electroweak penguin P_{EW} effects [7]. For the latter, NP CPV phase would be needed. Although the experimental significance for $\mathcal{A}_{K^+\pi^0} - \mathcal{A}_{K^+\pi^-}$ is relatively robust, at present it is not yet conclusive whether NP phase is absolutely called for.

A third possible hint, though not yet widely perceived, is the $\mathcal{S}_f$ value in $b \rightarrow c\bar{c}s$ decays itself. The Belle experiment gives [8] $\mathcal{S}_{J/\psi K^0} = 0.652 \pm 0.039 \pm 0.020$ using 386M $B\bar{B}$ pairs, which dominates the HFAG average of $\mathcal{S}_{c\bar{c}s} = 0.685 \pm 0.032$ [3] over charmonium modes. The latter is pulled up slightly by the BaBar result of $\mathcal{S}_{c\bar{c}s} = 0.722 \pm 0.040 \pm 0.023$ based on 227M $B\bar{B}$ pairs. Early Belle results based on a smaller data sample also averaged over many charmonium modes, and gave a higher value. It remains to be seen what would be the future update value, especially from BaBar.

$\mathcal{S}_{J/\psi K}$ or $\mathcal{S}_{c\bar{c}s}$ are traditionally identified with the CKM phase $\sin 2\phi_1/\beta$ as CPV phase in $B^0 - \bar{B}^0$ mixing, since $b \rightarrow c\bar{c}s$ decay should be dominantly tree level and has very little CPV phase. The issue is then in comparing direct measurement with the so-called “CKM” [9] or “UT” [10] fit results. These studies are able to fit to all data other than TCPV measurement in B decay, and “predict” the value for $\sin 2\phi_1/\beta = \mathcal{S}_{J/\psi K} (\cong \mathcal{S}_{c\bar{c}s})$ assuming 3 generation SM. Before the Belle 2005 result [8], these predictions tended to give $\sin 2\phi_1/\beta > 0.75$, suggesting a possible $\mathcal{S}_{J/\psi K} - \sin 2\phi_1/\beta|_{\text{fitter}}$ problem.

Is there something happening in $b \rightarrow c\bar{c}s$ decay, similar to what we may be seeing in $\Delta\mathcal{S}$ and $\mathcal{A}_{K^+\pi^0} - \mathcal{A}_{K^+\pi^-}$? Unfortunately this problem would be hard to pin down without adopting an explicit NP model and implementing it in the fitter. That is, the reference point of $\sin 2\phi_1/\beta|_{\text{fitter}}$ would itself become the point of con-

tention and is somewhat ill-defined.

Taking all three hints together, NP phases may well be present in the $b \rightarrow c\bar{c}s$ amplitude. Surprisingly, the way out may be DCPV in $B^+ \rightarrow J/\psi K^+$ mode, thanks to both its precision measurement nature, and the plausibility of an associated finite strong phase.

We aim to keep things as simple as possible. Let us put the $B \rightarrow J/\psi K$ decay amplitude in the form

$$\mathcal{M}(\bar{B} \rightarrow J/\psi \bar{K}) = a e^{i\delta} + b e^{-i\phi} = a(e^{i\delta} + \hat{b} e^{-i\phi}), \quad (2)$$

where the first term carrying the strong phase is defined with a positive and dominating the rate. We will maintain $a \cong \sqrt{\mathcal{B}(B \rightarrow J/\psi K)}$ throughout this note, treating the second term carrying the weak phase as a small correction to the rate, i.e. $\hat{b} \equiv b/a \ll 1$. From Eq. (2), to good approximation for small $\hat{b}$, one has

$$\mathcal{S}_{J/\psi K^0} \cong \frac{\sin 2\Phi_{B_d} + 2\hat{b} \sin(2\Phi_{B_d} + \phi) \cos \delta}{1 + 2\hat{b} \cos \phi \cos \delta}, \quad (3)$$

$$\mathcal{A}_{J/\psi K^+} \cong -2\hat{b} \sin \phi \sin \delta. \quad (4)$$

The CPV phase in B_d mixing is put in the more general form of Φ_{B_d} , which would be ϕ_1/β in SM. Let us analyze the strength of δ in general, as well as the strength $\hat{b}$ and phase ϕ within SM.

The $B \rightarrow J/\psi K$ decay is a color-suppressed process, and there is no factorization theorem. It is generated by the tree level effective a_2 coefficient, hence dominates a in Eq. (2). Since this effective a_2 is larger than naive, it contains hadronic effects that would likely generate a finite strong phase $\delta \cong \arg a_2$. For color-suppressed modes that are similarly enhanced, such as $B^0 \rightarrow D^0 \pi^0$, analysis [11] shows that the associated strong phase $\delta \sim 30^\circ$. Further evidence for such strong phase comes from experimental study of $B \rightarrow J/\psi K^*$ decay. By angular analysis, strong phase differences between the various helicity components vary from $\sim 26^\circ$ [12] to $\sim 34^\circ$ [13]. Although one is not directly measuring the strong phase of the longitudinal amplitude (the analogue of $B \rightarrow J/\psi K$), the generic strength is consistent with the $D^0 \pi^0$ result. Thus, we take $\delta \simeq 30^\circ$ as our nominal strong phase for $B \rightarrow J/\psi K$ decay.

Denoting Δ_{ct} and Δ_{ut} as the penguin loop amplitudes associated with the CKM coefficients $\lambda_c \equiv V_{cs}^* V_{cb}$ and $\lambda_u \equiv V_{us}^* V_{ub}$, in SM Eq. (2) becomes

$$\mathcal{M}_{\text{SM}} \simeq a \left[e^{i\delta} + \left| \frac{\lambda_u}{\lambda_c} \right| \frac{\hat{\Delta}_{ut}}{1 + \hat{\Delta}_{ct}} e^{-i\phi_3} \right], \quad (5)$$

where $\phi = \phi_3 = \arg V_{ub}^*$, and a now absorbs Δ_{ct} , but still saturates the decay rate. Note that to good approximation $\hat{\Delta}_{ij}$ is normalized by a . We see that

$$\hat{b}_{\text{SM}} \simeq \left| \frac{\lambda_u}{\lambda_c} \right| \frac{\hat{\Delta}_{ut}}{1 + \hat{\Delta}_{ct}}, \quad (6)$$

is rather small. Since $\lambda_c \Delta_{ct}$ should dominate $B \rightarrow \phi K$ decay [15], ignoring kinematic differences, we have the

crude estimate $|\hat{\Delta}_{ct}| \sim \sqrt{\mathcal{B}(B \rightarrow \phi K)/\mathcal{B}(B \rightarrow J/\psi K)} \sim 0.1$. Taking $|\lambda_u/\lambda_c| \simeq 0.02$ and saturating $|\hat{\Delta}_{ut}| \lesssim |\hat{\Delta}_{ct}|$, we get $\hat{b}_{\text{SM}} = \mathcal{O}(0.002)$, which implies a negligible shift in $\mathcal{S}_{J/\psi K^0}$. Together with $\phi \simeq \phi_3 \simeq 60^\circ$ and $|\sin \delta| \sim 1/2$, we find the order of magnitude estimate

$$\mathcal{A}_{J/\psi K^+}^{\text{SM}} = \mathcal{O}(0.003), \quad (7)$$

which can be improved by constraining $|\hat{\Delta}_{ut}/\hat{\Delta}_{ct}|$.

We now extend to consider New Physics scenarios. For its simplicity, we keep the form of Eq. (2) with a saturating the $B \rightarrow J/\psi K$ rate, and b the term carrying NP phase. In actual computations, the SM phase is kept in the numerics.

It should be clear from Eq. (3) that, assuming $\sin(2\Phi_{B_d} + \phi^{\text{new}})$ and $\cos \delta \sim 1$, to have $\mathcal{S}_{J/\psi K}$ shifted by ~ 0.05 or so, one needs $\hat{b}$ to be a couple of percent. From Eq. (4) we see that $\mathcal{A}_{J/\psi K}$ could then be a few percent as well. The question then is not whether a downward shift in $\mathcal{S}_{J/\psi K}$ could generate observable $\mathcal{A}_{J/\psi K}$, but whether this could be achieved in realistic models.

We will illustrate with New Physics in the electroweak penguin P_{EW} . Since P_{EW} arises from short distance, it should not have sizable strong phase, so assignment of strong phase to the a term in Eq. (2) is intuitive. We will consider two realistic models where New Physics CPV phase could appear in P_{EW} : the existence of a fourth generation, or an extra Z' .

In a four sequential generation model, natural for affecting P_{EW} , we have found [7] 1) $\mathcal{A}_{K^+\pi^0} - \mathcal{A}_{K^+\pi^-} \neq 0$ can be accounted for; 2) $b \rightarrow s\ell^+\ell^-$ rate and B_s mixing can be SM-like; 3) after satisfying kaon constraints, no visible deviation of Δm_{B_d} from SM, and $\Phi_{B_d} \sim \phi_1$ [14]; 4) $\Delta \mathcal{S}_{K^0\pi^0, \phi K^0} < 0$ can be generated, and is robust against hadronic uncertainties [4]. We now illustrate the new point 5): while $\sin 2\Phi_{B_d}$ mimics $\sin 2\phi_1/\beta$ (see Fig. 4 of Ref. [14]), $\mathcal{S}_{J/\psi K}$ is in fact driven lower to around 0.7, just at the right CPV phase value of $\phi_{sb} \simeq 70^\circ$ (defined by $V_{t's}^* V_{t'b} \equiv r_{sb} e^{i\phi_{sb}}$) for points 1)–4) above. We emphasize that point 2) has not changed with the measurement [16] of B_s mixing; a year ago we stated that [7] Δm_{B_s} should be SM-like in 4 generation model.

As discussed, we employ the SM-dominated parameter a to saturate the $B \rightarrow J/\psi K$ rate. The P_{EW} contribution b , arising from 4th generation and carrying CPV phase $e^{i\phi_{sb}}$, can be computed straightforwardly as for $B \rightarrow K\pi^0, \phi K$ modes. One essentially has $b \rightarrow s$ transitions induced by a virtual Z , which itself turns into π^0, ϕ , or here the J/ψ . We illustrate in Fig. 1(a) the $b \rightarrow d$ quadrangle. Defining $V_{t'd}^* V_{t'b} \equiv r_{db} e^{i\phi_{db}}$, the quadrangle with $V_{t'd}^* V_{t'b}$ in 3rd quadrant would lead to a high value of $\sin 2\Phi_{B_d} \simeq 0.78$, and is disfavored [14]. The quadrangle corresponding to $\phi_{db} \sim 10^\circ$ can hardly be distinguished from the triangle of the 3 generation case, hence mimics SM in $b \rightarrow d$ transitions. In Fig. 1(b) we plot $\mathcal{S}_{J/\psi K}$ vs ϕ_{sb} for our standard parameter set of $m_{t'} \sim 300$ GeV, $r_{sb} \sim 0.02, 0.025$ and 0.03 , with strong phase δ set to zero. $\mathcal{S}_{J/\psi K}$ can dip below 0.7 around $\phi_{sb} \sim 70^\circ$, for all




FIG. 1: (a) Unitarity quadrangle for $b \rightarrow d$ transitions with $V_{t'd}V_{t'b}^* \equiv r_{db}e^{-i\phi_{db}}$ exaggerated and horizontal scale shrunk by half, for $\phi_{db} \sim 10^\circ$ and 100° (dashed), compared with 3 generation SM (solid); (b) $S_{J/\psi K^0}$ vs ϕ_{sb} , where $V_{t's}^*V_{t'b} \equiv r_{sb}e^{i\phi_{sb}}$; (c) $S_{J/\psi K^0}$ and (d) $\mathcal{A}_{J/\psi K^+}$ vs δ , for $m_{t'} = 300$ GeV and $r_{sb} = 0.02, 0.025$ and 0.03 . For (b), we set $\delta = 0$. For (c) and (d), we set $\phi_{sb} = 70^\circ$. In (b), (c) and (d), $\phi_{db} = 10^\circ$, and larger r_{sb} gives stronger variation.

r_{sb} . Fig. 1(c) illustrates that a small strong phase δ of order $\pm 30^\circ$ or less does not affect this slight suppression of $S_{J/\psi K}$, but at 180° would almost restore it back. In Fig. 1(d) we plot $\mathcal{A}_{J/\psi K^+}$ vs δ . We see that $\mathcal{A}_{J/\psi K^+}$ can reach above 1% for δ up to $\pm 30^\circ$. This could become measurable soon.

To illustrate further the correlations between a lower $S_{J/\psi K^0}$ and a finite $\mathcal{A}_{J/\psi K^+}$, let us invoke a second model, that of extra Z' boson with flavor changing couplings. Z' models can also [17] explain the $\mathcal{A}_{K^+\pi^0} - \mathcal{A}_{K^-\pi^-}$ difference, as it can not only enter P_{EW} , but affect color-suppressed strong penguin amplitudes as well. In fact, Z' models are much less constrained than the previous 4th generation model, since one has freedom to assign the extra $U(1)$ couplings to various fermions. For instance, one can ask the Z' to be leptophobic, hence decouple from $b \rightarrow s\ell^+\ell^-$ constraint, while the consistency of measured Δm_{B_s} with SM expectation allows a range of Z' mass scales [18]. We therefore use the framework of Eqs. (2)–(4) again to illustrate first the possible impact on $S_{J/\psi K}$.

We follow the formalism introduced in Ref. [17], and adopt the particular choice for the Z' contributions to the Wilson coefficient at the weak scale. Assuming LL coupling, the weak phase is placed in the effective charge $\arg B_{sb} = \phi_L$, and the effective coupling is $\xi_{bsqq} = cB_{sb}B_{qq}$ for $b \rightarrow s\bar{q}q$ transition, where $c = (g'M_Z/gM_{Z'})^2/|V_{ts}^*V_{tb}|$, and B_{qq} is the effective qq charge. In $B \rightarrow K\pi$ case, for strong penguin one has $B_{qq} = B_{uu} + 2B_{dd}$, while for EW penguin one has $B_{qq} = 2(B_{uu} - B_{dd})$. By taking $B_{uu} = -2B_{dd}$ [17], the contribution would be only in P_{EW} . For our present




FIG. 2: (a) $S_{J/\psi K^0}$ vs ϕ_L for $\delta = 0$, and (b) $\mathcal{A}_{J/\psi K^+}$ vs δ for $\phi_L = -80^\circ$, for $(\xi_3^{LL}, \xi_9^{LL}) = (-0.001, -0.001)$ (dashed), $(-0.003, -0.003)$ (solid) and $(-0.005, -0.005)$ (dot-dashed), respectively.

$B \rightarrow J/\psi K$ case, we take $B_{cc} = B_{uu}$ and there is no B_{dd} contribution, we therefore parameterize

$$\Delta c_3^{\psi K}, \Delta c_9^{\psi K} = +\frac{2}{3}\xi_3^{LL}e^{i\phi_L} + \frac{4}{3}\xi_9^{LL}e^{i\phi_L}, \quad (8)$$

where $\Delta c_9^{\psi K}$ contributes to P_{EW} , while $\Delta c_3^{\psi K}$ is a color-suppressed strong penguin, and we take $\Delta c_5^{\psi K} = \Delta c_7^{\psi K} = 0$.

We plot $S_{J/\psi K^0}$ vs ϕ_L for $\delta = 0$ in Fig. 2(a), for three different combinations of the parameters $(\xi_3^{LL}, \xi_9^{LL}) = (-0.001, -0.001), (-0.003, -0.003)$ and $(-0.005, -0.005)$. We have checked that a small δ value does not affect $S_{J/\psi K^0}$, while the parameter choices do not violate current measurements of ΔS . We see that around $\phi_L \sim -90^\circ$ or so, $S_{J/\psi K^0}$ can be brought down below 0.7, and could even approach 0.65. Keeping $\phi_L \sim -80^\circ$, we plot $\mathcal{A}_{J/\psi K^+}$ vs δ in Fig. 2(b). We see that for $|\delta| \sim 30^\circ$, $|\mathcal{A}_{J/\psi K^+}|$ can be rather sizable, and is constrained in fact by Eq. (1). A value of a few % could be established with existing data, and by this summer with over 500M $B\bar{B}$ events at Belle alone.

We have not intended to go into detailed modelling, but just to illustrate that the possible hint for New Physics in a lower value of $S_{J/\psi K^0}$, could imply a direct CPV asymmetry $\mathcal{A}_{J/\psi K^+}$ at 1% level or higher. Our results imply that, if one takes the difference of $S_{J/\psi K^0} - \sin 2\phi_1/\beta^{\text{fit}}$ to be of order -0.05 , then $\mathcal{A}_{J/\psi K^+}$ could be as large as a few percent, as illustrated by the Z' model. This is consistent with the experimental bound of Eq. (1). Since the latter is based on only a fraction of the data that is currently at hand, there is much room for improvement.

We caution, however, that the strong phase δ , though of order 20° – 30° on general grounds, could be small *by accident*. So, $\mathcal{A}_{J/\psi K^+}$ could still be very small, even if large NP is present. In addition, for the more constrained 4th generation model, besides linking all the hints and giving more wide ranging predictions, the prediction for $\mathcal{A}_{J/\psi K^+}$ is at the more conservative 1% level. In this respect, it is desirable to give a better prediction for $\mathcal{A}_{J/\psi K^+}^{\text{SM}}$ than Eq. (7), by constraining the penguin amplitude ratio Δ_{ut}/Δ_{ct} for $B \rightarrow J/\psi K$ decay in SM.

Assuming 900M $B\bar{B}$ events between Belle and BaBar by this summer, the combined statistical error for

$\mathcal{A}_{J/\psi K^+}$ could reach below 0.005, and a 2% asymmetry, if present, should be clearly established. Assuming isospin symmetry, one could combine with $B^0 \rightarrow J/\psi K^0$, which requires tagging. But the error is reduced only by $\sim 10\%$ or so, unless time dependent analysis provides extra leverage. To probe $|\mathcal{A}_{J/\psi K^+}| \sim 1\%$, one needs to await the order 2000M events expected by 2008. With statistical error down to 0.003 or lower, systematic issues would become important. Studies on systematic errors at the per mill level should be very worthy towards the era of Super B factory, where many processes of interest would not be as clean as $B \rightarrow J/\psi K$.

We have focused on B factory prospects, but $B \rightarrow J/\psi K$ decay is accessible also at hadronic colliders. With 110 pb^{-1} data from Tevatron Run I, CDF reconstructed [19] 860 $B \rightarrow J/\psi K^+$ events with $J/\psi \rightarrow \mu^+ \mu^-$. Scaling to 8 fb^{-1} , one expects 6.2×10^4 events towards the end of Tevatron Run II. If the acceptance and efficiency can be improved by a factor of 2 or 3, the CDF and D0

experiments could be competitive with the $\sim 2 \times 10^5$ reconstructed $B \rightarrow J/\psi K^\pm$ events per 1000M $B\bar{B}$ pairs at the B factories. Systematics could be under better control at hadron colliders due to more abundant calibration modes. After 2008, with the advent of the LHC, the ATLAS, CMS and LHCb experiments should give the best measurement of $\mathcal{A}_{J/\psi K^+}$, perhaps even measuring the SM expectation at the few per mill level.

In summary, in view of possible hints of New Physics in CP violation in $b \rightarrow sq\bar{q}$ and $c\bar{c}s$ decays, we point out that the direct CP violation asymmetry $\mathcal{A}_{J/\psi K^+}$ could be at the 1% level or higher, and can be measured in the next few years. If established, this could become one of the most convincing evidence for New Physics.

Acknowledgement. This work is supported in part by NSC 94-2112-M-002-035 and NSC 94-2811-M-002-053 of Taiwan, and HPRN-CT-2002-00292 of Israel.

-
- [1] G. Bonvicini *et al.* [CLEO Collab.], Phys. Rev. Lett. **84**, 5940 (2000); K. Abe *et al.* [Belle Collab.], Phys. Rev. D **67**, 032003 (2003); B. Aubert *et al.* [BaBar Collab.], Phys. Rev. Lett. **92**, 241802 (2004).
 - [2] Particle Data Group, <http://pdg.lbl.gov/>.
 - [3] Heavy Flavor Averaging Group [HFAG], see webpage <http://www.slac.stanford.edu/xorg/hfag/>.
 - [4] W.S. Hou, M. Nagashima, G. Raz, A. Soddu, hep-ph/0603097, and references therein.
 - [5] B. Aubert *et al.* [BaBar Collab.], Phys. Rev. Lett. **93**, 131801 (2004); Y. Chao *et al.* [Belle Collab.], *ibid.* **93**, 191802 (2004); K. Abe *et al.* [Belle Collab.], hep-ex/0507045.
 - [6] See e.g. H.n. Li, S. Mishima and A.I. Sanda, Phys. Rev. D **72**, 114005 (2005), and references therein.
 - [7] W.S. Hou, M. Nagashima and A. Soddu, Phys. Rev. Lett. **95**, 141601 (2005); and references therein.
 - [8] K. Abe *et al.* [Belle Collab.], hep-ex/0507037.
 - [9] J. Charles *et al.* [CKMfitter Group], Eur. Phys. J. C **41**, 1 (2005); see <http://ckmfitter.in2p3.fr/> for updates.
 - [10] M. Bona *et al.* [UTfit Collab.], JHEP **0507**, 028 (2005); see <http://utfit.roma1.infn.it/> for updates.
 - [11] See e.g. C.K. Chua, W.S. Hou and K.C. Yang, Phys. Rev. D **65**, 096007 (2002).
 - [12] R. Itoh *et al.* [Belle Collab.], Phys. Rev. Lett. **95**, 091601 (2005).
 - [13] B. Aubert *et al.* [BaBar Collab.], Phys. Rev. D **71**, 032005 (2005).
 - [14] W.S. Hou, M. Nagashima and A. Soddu, Phys. Rev. D **72**, 115007 (2005).
 - [15] From $\mathcal{A}_{\phi K^+} = 0.037 \pm 0.050$ [3], with $a e^{i\delta} = \lambda_c \Delta_{ct}$ and $b e^{-i\phi} = \lambda_u \Delta_{ut}$, there seems to be little strong phase difference between Δ_{ct} and Δ_{ut} .
 - [16] G. Gomez-Ceballos [CDF Run II Collab.], talk at *Flavor Physics and CP Violation* Conference, April 2006, Vancouver, Canada.
 - [17] V. Barger, C.W. Chiang, P. Langacker, H.S. Lee, Phys. Lett. B **598**, 218 (2004).
 - [18] K. Cheung, C.W. Chiang, N.G. Deshpande, J. Jiang, hep-ph/0604223; P. Ball and R. Fleischer, hep-ph/0604249.
 - [19] F. Abe *et al.* [CDF Collab.], Phys. Rev. D **58**, 072001 (1998).